

## **General Disclaimer**

### **One or more of the Following Statements may affect this Document**

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

NASA TM X-65390

**RELATION OF THE PLASMAPAUSE  
POSITION TO THE REGION OF ENHANCED  
FLUXES OF TRAPPED ENERGETIC  
( $E > 280$  keV) ELECTRONS DURING  
THE JUNE 15, 1965 MAGNETIC STORM**

**D. L. CARPENTER  
C. G. PARK  
J. F. ARENS  
D. J. WILLIAMS**

**NOVEMBER 1970**



**GSFC**

**GODDARD SPACE FLIGHT CENTER  
GREENBELT, MARYLAND**

**N71-13025**

FACILITY FORM 602

(ACCESSION NUMBER)

15

(PAGES)

TMX 65390

(NASA CR OR TMX OR AD NUMBER)

(THRU)

G3

(CODE)

29

(CATEGORY)

RELATION OF THE PLASMAPAUSE POSITION TO THE REGION OF ENHANCED  
FLUXES OF TRAPPED ENERGETIC ( $E > 280$  keV)  
ELECTRONS DURING THE JUNE 15, 1965 MAGNETIC STORM

D. L. Carpenter and C. G. Park  
Radioscience Laboratory  
Stanford University  
Stanford, California

J. F. Arens  
NASA-Goddard Space Flight Center  
Greenbelt, Maryland 20771

and

D. J. Williams  
ESSA Research Laboratories  
Space Disturbance Laboratory  
Boulder, Colorado 80302

NOVEMBER 1970

RELATION OF THE PLASMAPAUSE POSITION TO THE REGION OF ENHANCED  
FLUXES OF TRAPPED ENERGETIC ( $E > 280$  keV)  
ELECTRONS DURING THE JUNE 15, 1965 MAGNETIC STORM

D. L. Carpenter and C. G. Park  
Radioscience Laboratory  
Stanford University  
Stanford, California

J. F. Arens  
NASA-Goddard Space Flight Center  
Greenbelt, Maryland 20771

and

D. J. Williams  
ESSA Research Laboratories  
Space Disturbance Laboratory  
Boulder, Colorado 80302

ABSTRACT

This is a report on the position of the plasmopause relative to the region of appearance of enhanced fluxes of energetic ( $E > 280$  keV) electrons during the magnetic storm of June 15, 1965. The plasmopause was measured near the prime geomagnetic meridian by whistler techniques; the trapped energetic electrons were detected by the satellite 1963-38C in polar orbit at 1100 km altitude. As the storm-time reduction in plasmasphere radius occurred, the electron fluxes increased in a region that was apparently exterior to the diminishing plasmasphere. The plasmopause position appears to have been adjacent to the region of inflation of the magnetosphere reported by Cahill from Explorer 26 magnetometer data. This relationship is similar to that deduced from the OGO-3 measurements of Frank (ring current particles) and Taylor and his colleagues (thermal ion density).

## Introduction

It is well known that the radius of the plasmopause is reduced during periods of increasing planetary magnetic disturbance (Carpenter, 1966, 1967; Taylor et al., 1965; Chappell et al., 1970a). For example, whistler observations during the active-sun years of 1958-1959 include several examples of plasmopause radii between 2 and 3  $R_E$  (Corcuff and Delaroche, 1964; Carpenter, 1963). Data from the August 16-19, 1959 magnetic storm indicate a minimum plasmopause radius of 2  $R_E$  or less (Carpenter, 1962).

Williams et al (1968) found that electrons with energy greater than several hundred keV are injected or energized deep within the magnetosphere during magnetic storms. These authors suggested that the energetic electrons initially appear exterior to the plasmopause. This note presents data on the plasmopause position and on the distribution of trapped energetic electrons ( $E > 280$  keV) during the early phases of the June 15, 1965 magnetic storm. As the storm-time reduction in plasmasphere radius occurs, the electron flux increases in a region that is apparently exterior to the diminishing plasmasphere, in agreement with the suggestion of Williams et al (1968).

## Sources of data

The whistler data on plasmopause radius were obtained from recordings at Eights, Antarctica at  $L \sim 4$  near the prime geomagnetic meridian. The energetic electron data, previously published by Williams et al. (1968), were obtained from the low-altitude ( $\sim 1100$  km) polar orbiting satellite 1963-38C. (The satellite is magnetically aligned and the detectors oriented to look normal to the alignment axis.) The results of the

comparison are summarized in Figure 1, which includes plots of the AE index,  $D_{st}$ , and  $K_p$  for the three-day period June 15-17, 1965. Arrows and triangles represent whistler information on plasmopause equatorial radius, while observations of energetic particles are shown by iso-intensity flux contours. In the case of the whistler data, an arrow means that the estimated plasmopause position is in the direction of the arrowhead with an uncertainty of about  $0.5 R_E$ . A filled triangle means that the position is in the direction of the apex and with uncertainty of about  $0.2 R_E$ . In the case of the trapped energetic electron data a set of open circles aligned vertically indicates where counts of 10, 100, or 1000/sec were observed on an individual satellite pass. The position of the peak intensity for the pass is indicated by an X. Dashed curves connect the circles to form crude iso-intensity flux contours, while the variation of the position of the peak with time is indicated by a solid curve.

The whistler data are essentially measurements of minimum B along field aligned paths. (For a review of the whistler method, see Angerami (1966), Angerami and Carpenter (1966), or Carpenter and Smith (1964) In the case of Figure 1, the values of minimum B were converted to equatorial radius by means of a dipole field model. Because of the shrinkage of the plasmasphere into a region of relatively large B during the storm main phase, estimated maximum corrections for ring current effects involve reducing dipole estimates by roughly  $0.1 R_E$ . (The estimated corrections are based on Cahill's magnetometer observations from Explorer 26 during the June, 1965 storm (Cahill, 1970).)

The particle data are plotted in terms of  $L$  calculated for the  $\sim 1100$  km altitude of the satellite in the main Jensen and Cain field (Williams et al., 1968). During the storm main phase, distension of the tubes of force was such that the corresponding equatorial  $L$  values should be somewhat larger than those indicated in the figure (Williams et al., 1968). As noted, there is an opposite effect in the whistler data; hence in terms of equatorial radius the plasmopause and curve of peak intensity should be displaced a few tenths of an earth radius further apart than is presently shown for late June 16 and early June 17.

A complication in the comparison of Figure 1 is that the whistler observations represent the thermal plasma as observed near the prime geomagnetic meridian, while the satellite data, involving high particle energies, represent an essentially worldwide picture (the satellite orbit was near the dawn-dusk meridian). The restriction of the whistler observations can be partially overcome through statistical information on plasmopause behavior during substorms and through results of a number of simultaneous plasmopause measurements that have been made at widely spaced longitudes. Some remarks on extrapolation of the whistler data to other longitudes will be made below.

#### Plasmopause-energetic electron comparison during June 15-17, 1965

Figure 1 shows an increase in AE activity to several hundred gamma following  $\sim 12$  UT on June 15. This increase is accompanied by an initial drop in  $D_{st}$  and a surge in  $K_p$ . As this activity develops the Eights meridian is moving into the morning sector (see upper, LT scale), and no immediate effect on the plasmopause radius is seen. This is consistent

with the known tendency for the dayside to be shielded from large perturbing substorm electric field (Mozer and Serlin, 1969; Carpenter, 1970). As the whistler observations move into the afternoon sector, substorm activity continues and there is a several-hour period during which measurable whistlers were not detected. When measurements could again be made near local dusk, about 00 UT on June 16, the estimated plasmopause radius was lower than before, although details remain uncertain because of an absence of clear whistler traces propagating outside the plasmopause. Pronounced reduction in apparent plasmopause position continues near and after midnight local time on the 16th as the AE index surges toward a very high maximum and the  $D_{st}$  level drops rapidly. By local dawn on June 16 the plasmopause position is well defined near  $2.4 R_E$  (filled triangles). This reduction was achieved near the time of maximum substorm activity, and is consistent with the type of nightside main-phase behavior observed during several less severe magnetic storms in 1963, when the plasmopause radius near dawn was reduced to about  $3 R_E$  (Carpenter, 1966).

Following the minimum in plasmopause radius there is an outward trend on the local dayside of the 16th, apparently to about  $3 R_E$ . Details are not clear, but the information now comes from the outer, low-density region and thus provides information on the outer, rather than inner, limit of the plasmopause position. The minimum in  $D_{st}$  occurs at roughly 00 UT on the 17th, and by the end of the 17th the more rapid part of the recovery of  $D_{st}$  is completed.



On June 17, AE activity is high but shows a pronounced quieting trend. Across local nightside on the 16-17th the plasmopause again shows an inward trend, less dramatic than on the 15-16th. (Some details of this inward trend are described by Carpenter et al. (1969b).)

The iso-intensity contours in Figure 1 show that fluxes of electrons with  $E > 280$  keV began to build up rapidly between 4 and 5  $R_E$  following 00 UT on the 16th. The region of apparent injection is relatively broad, extending from 3.5 to about 5  $R_E$ . In this region the fluxes reach half their maximum value in about half a day, while fluxes at higher and particularly at lower L values require longer, thus giving the impression of an injection near  $L = 4.5$  and subsequent diffusion to higher and lower magnetic shells. Figure 1 indicates that the bulk of the "injection" event takes place in the region beyond the plasmopause, insofar as the latter is defined from the Eights meridian. The figure also indicates some overlap of the plasmasphere and energetic electrons on June 17 following a period of rapid inward diffusion of the electrons late on the 16th.

Following June 17 the energetic electrons continued to diffuse to lower and higher L shells in the manner reported by Williams et al. (1968). The comparison of Figure 1 is terminated at the end of June 17 due to complexities in describing the plasmopause position during the latter recovery phase of a storm. The recovery process is often complex, involving an interplay between continued substorm agitation and the slow filling of tubes of ionization outside the main-phase position of the plasmopause (Park, 1970). During quieting, the conditions for the

establishment of the plasmopause may exist at high L values before a steep gradient in the thermal plasma is readily detectable there, and while the thermal plasma at lower L shells retains an imprint of the storm-time plasmopause (see, for example, Chappell et al., (1970b); Carpenter (1970)).

### Discussion

It is possible to make a crude estimate of the world-wide behavior of the plasmopause by use of statistics on storm-time behavior near the Eights meridian in 1963 and 1965 (cf. Carpenter and Stone (1968); Carpenter (1970)). From the intensity and long duration of substorm activity on June 15-16, it is inferred that the entire plasmasphere was significantly reduced in size during the early phase of the storm. From the above-mentioned shielding of the dayside against substorm convection effects and the associated concentration of erosion and/or compression effects in the dusk, dawn, and midnight sectors, it is conjectured that the Eights meridian was nearly the last to exhibit a reduction in plasmopause radius. Thus the plasmopause data for other meridians would probably resemble the data of Figure 1, but would be shifted to the left by varying amounts. As activity developed on the latter half of the 15th UT, regions well to the west of Eights would have been in the midnight-dawn sector, and would have experienced immediate inward displacements, probably to within  $4 R_E$ . Regions well to the east of Eights would have reached the dusk meridian near say 18 UT on the 15th, and hence would have begun a cycle of erosion and compression events at an earlier time. Thus it is probable that by 00 UT on the 16th, the plasmopause radius was everywhere less than about  $4 R_E$ .

Direct evidence of the worldwide shrinkage of the plasmasphere was provided on June 17 at  $\sim 10$  UT by simultaneous measurements from OGO 1 and from Eights (Carpenter et al., 1969a). Taylor's ion mass spectrometer on OGO 1 showed a plasmopause crossing at  $L \sim 3.5$  near 13 LT, while the simultaneous Eights whistler data (Figure 1) indicate a radius of  $\sim 3 R_E$  at  $\sim 05$  LT. This is good agreement when expected variations with longitude and local time are taken into account.

From the results summarized in Figure 1 and from the foregoing discussion, it is tentatively concluded that the 'injection' event took place outside the plasmasphere, in agreement with the earlier suggestion of Williams et al (1968). Energetic electrons later appeared within the outer plasmasphere, following what was apparently a period of rapid cross-L diffusion.

From Explorer 26 magnetometer data Cahill (1970) has shown that the asymmetric, main phase of the June storm involved magnetospheric inflation concentrated in the L range  $\sim 2.7 - 3.7 R_E$ . This L range flanks and possibly somewhat overlaps the plasmopause as observed from Eights on the 16th of June. Assuming a worldwide reduction in plasmopause radius roughly comparable to that observed from Eights, this close spatial relation of ring current and plasmopause appears similar to that reported by Frank (1967) and Taylor et al. (1968) from observations during the July 9, 1966 storm.

ACKNOWLEDGMENTS

The research at Stanford was supported in part by the Office of Polar Programs of the National Science Foundation under grants GA-1151 and GA-1485, in part by the Atmospheric Sciences Section of the National Science Foundation under grants GA-775, GA-1445 and GA-10719 and in part by the National Aeronautics and Space Administration under grant ~~NGL-020-008~~. NGL 05-020-008

## REFERENCES

- Angerami, J. J., A whistler study of the distribution of thermal electrons in the magnetosphere, SEL-66-017 Radioscience Lab., Stanford Electronics Labs., Stanford University, Stanford, Calif., May 1966.
- Angerami, J. J. and D. L. Carpenter, Whistler studies of the plasmopause in the magnetosphere-2; equatorial density and total tube electron content near the knee in magnetospheric ionization, J. Geophys. Res., 71, 711, 1966.
- Cahill, L. J., Jr., Magnetosphere inflation during four magnetic storms in 1965, J. Geophys. Res., 75, 3778, 1970.
- Carpenter, D. L., New experimental evidence of the effect of magnetic storms on the magnetosphere, J. Geophys. Res., 67, 135, 1962.
- Carpenter, D. L., Whistler evidence of a 'knee' in the magnetospheric ionization density profile, J. Geophys. Res., 68, 1675, 1963.
- Carpenter, D. L., Whistler studies of the plasmopause in the magnetosphere -1; temporal variations in the position of the knee and some evidence on plasma motions near the knee, J. Geophys. Res., 71, 693, 1966.
- Carpenter, D. L., Relations between the dawn minimum in the equatorial radius of the plasmopause and  $D_{st}$ ,  $K_p$ , and local  $K$  at Byrd Station, J. Geophys. Res., 72, 2969, 1967.
- Carpenter, D. L., Whistler evidence of the dynamic behavior of the dusk-side bulge in the plasmasphere, J. Geophys. Res., 75, 3837, 1970.
- Carpenter, D. L., C. G. Park, H. A. Taylor, Jr., and H. C. Brinton, Multi-experiment detection of the plasmopause from EOGO satellites and Antarctic ground stations, J. Geophys. Res., 74, 1837, 1969a.

- Carpenter, D. L. and R. L. Smith, Whistler measurements of electron density in the magnetosphere, Rev. of Geophys., 2, 415, 1964.
- Carpenter, D. L. and K. Stone, Recent whistler research on hydromagnetic motions in the plasmasphere, Paper presented at International Symposium on the Physics of the Magnetosphere, Washington, D. C., Sept. 1968.
- Carpenter, D. L., K. Stone and S. Lasch, A case of artificial triggering of VLF magnetospheric noise during the drift of a whistler duct across magnetic shells, J. Geophys. Res., 74, 1848, 1969b.
- Chappell, C. R., K. K. Harris and G. W. Sharp, A study of the influence of magnetic activity on the location of the plasmopause as measured by OGO 5, J. Geophys. Res., 75, 50, 1970a.
- Chappell, C. R., K. K. Harris and G. W. Sharp, The morphology of the bulge region of the plasmasphere, J. Geophys. Res., 75, 3848, 1970b.
- Corcuff, Y. and M. Delaroche, Augmentation du gradient d'ionisation dans la proche magnétosphère en périodes de forte activité magnétique, C. R. Acad. Sci. Paris, 258, 650, 1964.
- Frank, L. A., On the extraterrestrial ring current during geomagnetic storms, J. Geophys. Res., 72, 3753, 1967.
- Mozer, F. S. and R. Serlin, Magnetospheric electric field measurements with balloons, J. Geophys. Res., 74, 4739, 1969.
- Park, C. G., Whistler observations of the interchange of ionization between the ionosphere and the protonosphere, J. Geophys. Res., 75, 4249, 1970.
- Taylor, H. A., Jr., H. C. Brinton and M. W. Pharo, III, Contraction of the plasmasphere during geomagnetically disturbed periods, J. Geophys. Res., 73, 961, 1968.

Taylor, H. A., Jr., H. C. Brinton, and C. R. Smith, Positive ion composition in the magnetosphere obtained from the OGO-A satellite, J. Geophys. Res., 70, 5769, 1965.

Williams, D. J., J. F. Arens and L. J. Lanzerotti, Observations of trapped electrons at low and high altitudes, J. Geophys. Res., 73, 5673, 1968.

## FIGURE CAPTIONS

Figure 1. Above. Comparison of the plasmopause position (arrows and triangles) and observations of trapped energetic electrons with  $E > 280$  keV (dashed flux contours) for June 15-17, 1965. There is a buildup or 'injection' of energetic electrons in a region apparently exterior to the diminishing plasmasphere. Equatorial radius is plotted vs universal time (bottom) and local time of the plasmopause observations (top). The plasmopause position was estimated from whistlers recorded at Eights, Antarctica ( $L \sim 4$ ) near the prime geomagnetic meridian. The trapped electron data is from satellite 1963-38C in polar orbit at  $\sim 1100$  km. See text for further details. Below. Plots of the hourly auroral electrojet index,  $D_{st}$ , and  $K_p$ .



